INDEX

Acolus mellilus, life history of, 816
Aenasius paulistus, 201
Aerial dusting, 912
  spraying, 676
Agromyza angelicae, 199
Agromyzid fly infesting sweetpotato seed, 704
Alabama argillacea, 830
Alfalfa snout beetle
  baits for control of, 639
  control of, 769
  fumigation of soil, 460
Alfalfa weevil, 892
  one new Arizona, 119
Amazon fly, 738
Ambrosia beetle transmits Dutch elm disease, 819
Anabasine, use against apple aphids, 800
Anacampsis fragariella, 121
Anasa tristis, 475
  parasitic castration of, 269
Anthonomus grandis, 682
Anticarsia gemmatilis, 635
Ants, fire, damage, 700
Anuraphis roseus, 336
Aonidiella aurantii, 895
Aphelinus mali, natural dispersion of, 811
Aphid injury and vetch varieties, 833
Aphids
  and narcissus mosaic, 942
  South Sea Island, 200
Aphis
  maidis, susceptibility of corn hybrids to, 623
  rumicis, toxicity of active nicotines to, 734
Apple
  aphids
    technique in experiments, 443
    tests of insecticides against, 800
  delayed dormant spray, 440
  maggots, methyl bromide against, 817
  mealy bug on, 445
  spray deposits, 505
Apple trees, spraying with semi-refined oils, 332
Arsenical grasshopper poisons, effect upon pheasants, 501
Arsenate sprays with soybean flour for codling moth, 862
Artichoke plume moth, 66
Asiatic beetle grubs, control with methyl bromide, 486
Aspidiotus perniciosus, 890
Attractants, for codling moth, 933
  larvae, 342

Bacillus larvae
  nitrite nitrogen test for, 405
  thermal resistance of, 399
Bait, poison, for strawberry crown borer, 812
Baits
  bacteria in, attracting codling moths, 933
  for firebrats, 857
  Japanese beetle, 327
  new, for alfalfa snout beetle, 639
  poison, for grasshopper control, 745
Barium compounds against firebrats, 857
Bathyplectes curculionis, 892
Bean weevils, destroyed by *Pediculoides ventricosus*, 581
Bee
  foods, vitamin content, 396
  losses in Utah, 418
  stings, reaction to, 418
Bees, package, 77
Beet
  leafhopper on tomatoes repelled by pyrethrin dusts, 389
  webworm, Hawaiian, control of, 272
Benzene against tobacco flea beetle, 834
Big-eyed bug bites man, 420
Biological control of
  codling and oriental fruit moths, 431
  mealybugs, 754
Black
  blister beetle, effect of pyrethrum upon, 807
  carpet beetle, effect of paradichlorobenzene upon, 816
  peach aphis, control, 204
  scale,
    frequency distribution of, 113
    hydrocyanic acid against eggs, 107
    winter mortality, 534
  widow spiders, ingestion with canned food, 550
Blattella germanica, 610
Blissus leucopterus, 688
Blow-fly
  larvae, method of destroying in soil, 662
  populations, 166
Blueberry bud mite, 699
Boll weevil
  control, 682, 787
  insecticides, effect on aphids and mirids, 792
Brachyrhinus ligustici, 460, 639
  control of, 769
Bruchidae in cowpea seed pods, 844
Bruchus pisorum, 881
Bud mite, new, on blueberry, 699
Budmoth, field tests for, 336
Bumblebees, insects in nests of, 81

Cabbage
  caterpillars, control, 463
  insects, control by derris-nicotine dusts, 416
Calcium
  arsenate, preparation of, 159
  arsenates, for boll weevil control, 682
  cyanamid, against pear thrips, 950
  cyanide, in control of squash bug, 575
Carnations, infested by onion thrips, 926
Carpet beetles, traps for control of, 651
Carpocapsa pomonella, 143, 340
Catalpa, mealy bug on, 445
Chalepus dorsalis, a pest of soybean, 742
Cherry fruit fly, control, 447
Chicken lice, effect of phenothiazine on, 700
Chinch bug, effect of fertilizers, 688
Chinch bug, hairy, control of, 153
Chloropicrin, fumigation of grain with, 536
Chloropids in houses, 946
Cirphis unipuncta, in Mississippi, 768
Citrus
  potato leafhopper on, 83
  rust mite, cost of control, 393
  tree spraying, rating thoroughness, 85
Clothes moth larvae, openings for, 582
Clothes moths, traps for control of, 651
Coal tar insecticides, new, 596
Coccinellid, introduced into California, 202
Cochliomyia americana, 622, 847, 850
Cockroaches, toxicity of phosphorus to, 193
Codling moth,
  activity of adults, 162
  and organic insecticides, 909

INDEX

Acolus mellilus, life history of, 816
Aenasius paulistus, 201
Aerial dusting, 912
  spraying, 676
Agromyza angelicae, 199
Agromyzid fly infesting sweetpotato seed, 704
Alabama argillacea, 830
Alfalfa snout beetle
  baits for control of, 639
  control of, 769
  fumigation of soil, 460
Alfalfa weevil, 892
  one new Arizona, 119
Amazon fly, 738
Ambrosia beetle transmits Dutch elm disease, 819
Anabasine, use against apple aphids, 800
Anacampsis fragariella, 121
Anasa tristis, 475
  parasitic castration of, 269
Anthonomus grandis, 682
Anticarsia gemmatilis, 635
Ants, fire, damage, 700
Anuraphis roseus, 336
Aonidiella aurantii, 895
Aphelinus mali, natural dispersion of, 811
Aphid injury and vetch varieties, 833
Aphids
  and narcissus mosaic, 942
  South Sea Island, 200
Aphis
  maidis, susceptibility of corn hybrids to, 623
  rumicis, toxicity of active nicotines to, 734
Apple
  aphids
    technique in experiments, 443
    tests of insecticides against, 800
  delayed dormant spray, 440
  maggots, methyl bromide against, 817
  mealy bug on, 445
  spray deposits, 505
Apple trees, spraying with semi-refined oils, 332
Arsenical grasshopper poisons, effect upon pheasants, 501
Arsenate sprays with soybean flour for codling moth, 862
Artichoke plume moth, 66
Asiatic beetle grubs, control with methyl bromide, 486
Aspidiotus perniciosus, 890
Attractants, for codling moth, 933
  larvae, 342

Bacillus larvae
  nitrite nitrogen test for, 405
  thermal resistance of, 399
Bait, poison, for strawberry crown borer, 812
Baits
  bacteria in, attracting codling moths, 933
  for firebrats, 857
  Japanese beetle, 327
  new, for alfalfa snout beetle, 639
  poison, for grasshopper control, 745
Barium compounds against firebrats, 857
Bathyplectes curculionis, 892
Bean weevils, destroyed by *Pediculoides ventricosus*, 581
Bee
  foods, vitamin content, 396
  losses in Utah, 418
  stings, reaction to, 418
Bees, package, 77
Beet
  leafhopper on tomatoes repelled by pyrethrin dusts, 389
  webworm, Hawaiian, control of, 272
Benzene against tobacco flea beetle, 834
Big-eyed bug bites man, 420
Biological control of
  codling and oriental fruit moths, 431
  mealybugs, 754
Black
  blister beetle, effect of pyrethrum upon, 807
  carpet beetle, effect of paradichlorobenzene upon, 816
  peach aphis, control, 204
  scale,
    frequency distribution of, 113
    hydrocyanic acid against eggs, 107
    winter mortality, 534
  widow spiders, ingestion with canned food, 550
Blattella germanica, 610
Blissus leucopterus, 688
Blow-fly
  larvae, method of destroying in soil, 662
  populations, 166
Blueberry bud mite, 699
Boll weevil
  control, 682, 787
  insecticides, effect on aphids and mirids, 792
Brachyrhinus ligustici, 460, 639
  control of, 769
Bruchidae in cowpea seed pods, 844
Bruchus pisorum, 881
Bud mite, new, on blueberry, 699
Budmoth, field tests for, 336
Bumblebees, insects in nests of, 81

Cabbage
  caterpillars, control, 463
  insects, control by derris-nicotine dusts, 416
Calcium
  arsenate, preparation of, 159
  arsenates, for boll weevil control, 682
  cyanamid, against pear thrips, 950
  cyanide, in control of squash bug, 575
Carnations, infested by onion thrips, 926
Carpet beetles, traps for control of, 651
Carpocapsa pomonella, 143, 340
Catalpa, mealy bug on, 445
Chalepus dorsalis, a pest of soybean, 742
Cherry fruit fly, control, 447
Chicken lice, effect of phenothiazine on, 700
Chinch bug, effect of fertilizers, 688
Chinch bug, hairy, control of, 153
Chloropicrin, fumigation of grain with, 536
Chloropids in houses, 946
Cirphis unipuncta, in Mississippi, 768
Citrus
  potato leafhopper on, 83
  rust mite, cost of control, 393
  tree spraying, rating thoroughness, 85
Clothes moth larvae, openings for, 582
Clothes moths, traps for control of, 651
Coal tar insecticides, new, 596
Coccinellid, introduced into California, 202
Cochliomyia americana, 622, 847, 850
Cockroaches, toxicity of phosphorus to, 193
Codling moth,
  activity of adults, 162
  and organic insecticides, 909

attracted by bacteria in baits, 933
biological control of, 431
control by arsenates and soybean flour, 862
effect of road dust upon, 803
flight habits, 436
larval attractants, 342
new habits of, 200
non-residue sprays for, 340
on stone fruits, 944
pyrethrum against, 143
sprays, 713
spur-burrowing habit of larvae, 796
Color, and effectiveness of Japanese beetle traps, 320
Comstock's mealybug on apple and catalpa, 445
Contributions, scientific at port of New York, 374
Corn borer, mechanical husker in control of, 643
Corn ear worm
control, by dichloroethyl ether, 584
metabolism in, 282
hibernates in Washington, 951
Corn hybrids, susceptibility to *Aphis maidis*, 623
Cotton
flea hopper, host plants, 456
insects, tests of insecticides for, 684
leafworm, 830
Cowpea seed pod insects
Crambus spp., 886
Cresylic acid, replaced with tar oil, 440
Cryolite insecticides, 699
Cubé dusts for pea weevil, 881
Curculionidæ in cowpea seed pods, 844
Cutworm, method for production of, 462
Cyclocephala immaculata, as a test insect, 704
Cylas formicarius, 921

Dendroctonus frontalis, 554
Deposits, mineral oil, on plant surfaces, 155
Derris
dust mixtures, peanut oil in, 736
dusts for pea weevil, 881
effect on black blister beetle, 807
-nicotine dusts for cabbage insects, 416
smoke, toxicity to housefly, 329
Dichloroethyl ether
for gladiolus thrips, 417
for pear thrips, 703
for wireworm control, 280
in mineral oil, against corn earworm, 584
Dinitro-o-cyclohexylphenol, in dormant sprays, 385
Dipping vat, for sheep parasites, 547
Disease, milky, of Japanese beetle, 303
DN dusts, for control of red spider on hops, 614
Dormant spray, for apples, 440
Dust, road, effect on codling moth, 803
Dusting
tent, 942
turf, new machine for, 153
Dusts, insecticidal, application of, 912
Dutch elm disease
and European elm bark beetle, 556
transmitted by *Scolytus sulcatus*, 250

Earwig, European, in Utah, 204
Elm leaf beetle,
control, 264
parasite, biology of, 947
Empoasca fabae, 83
Entomological
considerations, in use of insect killed pine, 773
observations, in California, 141
progress, fifty years of, 8, 15, 21, 30, 58
relations, Federal-State, 779, 784
Entomology,
development of research in, 749
Epicauta pennsylvanica, effect of pyrethrum and derris upon, 807
Epitrix parvula, 467, 834
Equipment, new, for heating honey, 410
Eriophyes vaccinii, 699
Erynnia nitida, 947
Ethylene dichloride, on peach, 199
Eufallia unicostata, new to Puerto Rico, 810
Eugenol and geraniol, attractiveness for Japanese beetle, 818
Euphasiopteryx australis, 202
European
earwig, in Utah, 204
elm bark beetle and Dutch elm disease, 556
European
pine shoot moth, 941
red mite, control of, 385

Federal-State relations, 779, 784
Filbert worm, 852
Fire
ants, damage, 700
-brat baits, 857
-weed nectar and humidity, 870
Fleas, economic importance, 701
Food habits of sparrow hawks, 583
Forficula auricularia, 204
Fumigants, naphthalene and nicotine, 368
Fumigation
of grain with chloropicrin, 536
soil, for alfalfa snout beetle, 460
tests in open tobacco warehouses, 812
with methyl bromide, 176, 921
Fungous disease, epidemics of, 201

Galerucella xanthomelaena, 264, 947
Gasterophilus nasalis, habits of larvae, 382
Geocoris decoratus, bite of, 420
Geraniol and eugenol, for Japanese beetle, 818
Geraniol, properties of, 327
Gladiolus thrips, control, 417, 490
Grape plume moth, 372
Grapholitha molesta, 189, 346
Graphs, for time-dosage mortality data, 139
Grasshopper
control by poison baits, 745
populations, effect of idle land upon, 646
Grasshoppers, a factor in soil erosion, 498
Green bug
injury, survey by airplane, 762
1939 outbreak in Oklahoma, 628
Golden codling moth, 582
Gulf coast tick, biology of, 179
Gypsy moth control, value to uninfested states, 558

Hairy chinch bug control, 153
Hawaiian beet webworm, control of, 272
Heliothis
armigera
hibernation in Washington, 951
metabolism in, 282
obsoleta, 129
Hessian fly, inheritance of resistance, 580
Hive temperature, effect of color of hive covers on, 646
Hollycock, a new pest of, 199
Holly leaf miner, 948
Honey, new equipment for heating, 410
Honeybee
combs,

infested by Mediterranean flour moth, 949
pests of, 81
pollen, trapping, 866
toxicity of tartar emetic to, 872
Hop
insects, field experiments, 112
pests, in Oregon, 70
Hops, control of red spider on, 614
Housefly, control by derris and pyrethrum, 329
Hydrocyanic acid
against red scale, 895
toxicity to eggs of black scale, 107
Hymenia fascialis, 272
Hypera brunneipennis, 119
postica, 892

Illinoia pisi, 833
Insect
activity at light traps, 654
aliens, undesirable, 1
control, effect of wheat harvesting machinery on, 646
damage, evaluation of, 499
food of trout, 544
-killed Ponderosa pine, 773
metabolism and temperature, 361
problems and changes in agriculture, 493
vectors and plant viruses, 827
Insecticidal
dusts, application of, 912
effects of organic compounds, 875
properties of *Phellodendron* spp., 941
residues in cured tobacco, 819
Insecticide
deposits, effect of weather, etc., on, 505
tests against pecan nut casebearer, 379
Insecticides
cryolite, 699
effect of upon aphids and mirids, 792
for boll weevil control, 787
for tobacco flea beetle, 834
household, effect on German cockroach, 610
Insecticides
new coal tar, 596
new, for shade trees, 247
organic, for codling moth, 909
rotenone bearing, 264
Insects
affecting combs of honeybee, 81, 949
cotton, 684
dried fruit, 115
entomophagous, 814
hop, 70
intercepted at New York, 374
list of common names, 708
living, for classroom, 123
of cowpea seed pods, 844
sucking, injuring rose, 658

Japanese beetle
attractiveness of geraniol and eugenol, 818
control by methyl bromide fumigation, 296
milky disease of, 303, 306
traps, 309, 317

Lead arsenate
commercial, particle size of, 170
residues from, 918
substitutes, 447
Leafhoppers, reaction to lawn sprinkling, 584
Leafmining sawfly on violet, 951
Leaf roller, control on roses, 364
Lepidoptera in cowpea seed pods, 844
Lepinotus patruelis, 946
Leptothrips mali, 539
Lice, chicken, phenothiazine on, 700
Light traps for
leafhoppers, 701
mosquitoes, 578
Lime sulfur, effect on
San Jose scale females, 890
treehopper eggs, 583
Locust leaf miner injuring soybeans, 742
Lygaeus kalmii, toxicity of nitrogenous bases to eggs of, 724

Macrosiphum pisi, 620, 736
Male scale, on olives, 203
Mealybug, biological control of, 754
Mediterranean flour moth, in honeybee combs, 949
Melissopus latiferreanus as pest of filberts, 852
Melonworm, control by rotenone dusts, 840
Metagonistylum minense, a parasite of sugarcane borer, 738
Methyl bromide
against
apple maggots, 817
artichoke plume moth, 66
Asiatic beetle, 486
pod borer larvae, 176
sweetpotato weevil, 921
fumigation for Japanese beetle, 296
in aqueous solution, 531
Mexican bean beetle, paris green as repellent, 525
Midges, aquatic, control and biology of, 252
Mineral oil deposits, determination on plants, 155
Milky disease of Japanese beetle larvae, 303
Milky diseases and Japanese beetle larvae, 306
Mosquito
collections,
by light traps, 578, 944
larvae
rearing of, 172
resistance to rotenone, etc., 172
repellents, studies of, 563, 567

Naphthalene fumigants, toxicity of, 368
Narcissus mosaic and aphids, 942
Nectar
in firewood, 870
secretion in poinsettia blossoms, 409
Nicotine
fumigants, toxicity of, 368
peat, sprays for use with, 717
persistence of, on rose bushes, 722
resistance of mosquito larvae to, 172
sulfate, combinations for rose bushes, 722
toxicity of internal doses, 511
use against apple aphids, 800
Nicotines
toxicity of, 734
Nitrogenous bases, toxicity of to eggs of *Lygaeus kalmii*, 724

Oeciacus vicarius, attacking man, 949
Oil
deposit, determination of by color, 905
spray residue, determination of, 900
sprays, effect on treehopper eggs, 583
toxicants and solids added to, 92
Oils, semi-refined for summer spraying, 332
Olives, maple scale on, 203
Oncideres twig girdlers, 561

Onion thrips
control, 278
sprays against, 926
Organic
compounds
insecticidal effects of, 875
toxicity of, to screwworms, 669
insecticides, status of for codling moth, 909
Oriental fruit moth
biological control of, 431
in Japan, 189
larvae parasitization of, 349, 353
Oxyptilus periscelidactylus, 372

Pantomorus
leucoloma, control by methyl bromide, 531
peregrinus, control by methyl bromide, 531
Paradichlorobenzene
against tobacco flea beetle,
effect on black carpet beetle, 816
on peach, 199
Paris green, particle size, 525
Parasite of
Pseudococcus maritimus, 201
Puerto Rican changa, 202
Parasitic castration of *Anasa tristis*, 269
Parasites
external of sheep, 547
of oriental fruit moth, 346, 349, 353
Pea
aphid, 736
as influenced by alfalfa plantings, 137
control, 72
in Maryland, 477
in Virginia, 620
rate of reproduction, 482
weevil control, 881
Peach
control of plum curculio, 451, 453
trees, mounding of, 199
Peanut oil in derris dust mixtures, 736
Pear thrips
control by dichlorethyl ether, 703
controlled by calcium cyanamide, 950
Peat, nicotine, sprays for use with, 717
Peacan nut casebearer, tests of insecticides, 379
Pediculoides ventricosus, destroying bean weevils, 581
Pentatomids, control on tomatoes, 275
Pest control, uniformity in, 776
Petroleum oil distillates, properties of, 591
Phellodendron spp., insecticidal properties of, 941
Phenothiazine,
effect on chicken lice, 700
toxicity to screwworms, 666
Phosphorus, against cockroaches, 193
Phyllophaga
crinita, life history of, 572
lanceolata, life history of, 572
Phytomyza ilicis, 948
Phytotoxic properties, of petroleum oil distillates, 591
Pickleworm, control by rotenone dusts, 840
Pine Shoot Moth, European, in British Columbia, 941
Platynota flavedana, 364
Platyptila carduidactyla, 66
Plot arrangement for timing applications, 357
Plum curculio
control, 451
in Georgia, 453
survival, 950
Pod borer larvae, methyl bromide against, 176
Poinsettia blossoms, nectar secretion in, 409
Poisons, stomach, effect of temperature on, 760
Pollen
substitutes, value of, 397
value of, 397
yields, 866
Popillia japonica, 296
Potato leafhopper
control on citrus, 83
Pseudococcus
comstocki, 445
longispinus, 754
maritimus, parasite of, 201
Psocids infesting
seed, 946
Pyrethrin dusts, repellency of, 389
Pyrethrum
effect on black blister beetle, 807
for control of codling moth, 143
smoke, toxicity to housefly, 329

Quarantine, and insect vectors, 827

Red scale
control, 92
by hydrocyanic acid, 895
Red spider
on hops, control of, 614
tartar emetic for, 579
Repellents, mosquito, value of, 563, 567
Residue
lead and arsenic ratio in, 918
oil spray, determination of, 900
Rhagoletis cingulata, 447
Rose
injury to by sucking insects, 658
leaf roller, control, 364
Rosy aphid, field tests for, 336
Rotenoid, definition of, 416
Rotenone
against apple aphids, 800
-bearing insecticides and the elm leaf beetle, 264
dusts for pickle and melon worms, 840
resistance of mosquito larvae to, 172

Saissetia oleae, 107
winter mortality, 534
San Jose scale, effect of lime-sulfur on females, 890
male, on olive, 203
Sawfly, leafmining, on violet, 951
Scolytus
multistriatus, mating habits, 818
sulcatus, 250
and Dutch elm disease, 250
Screwworm
larvae, nutrition of, 850
migration and pupation, 847
Screwworms, toxicity of
organic compounds to, 669
phenothiazine to, 666
Seed
-infesting psocid, 946
sweetpotato infested by Agromyzid fly, 704
Selenium
as toxicant for insects, 517
for plant injections, 517
Shade trees, insecticides for, 247
Sheep parasites, use of portable vat, 547
Simuliids, feeding activity of, 915
Sod webworms, biology and control, 886
Southern pine beetle, control of, 554
Soybean flour in codling moth control, 862

Soybeans, injured by locust leaf miner, 742
Sparrow hawks, food habits of, 583
Spilonota occellana, 336
Spraying
  aerial, with concentrated mixtures, 676
  citrus trees, 85
Spray
  materials, for use with nicotine peat, 717
  oil, toxicants and solids added to, 92
  residues, 447, 918
    oil, 900
Sprays
  codling moth, 713
  factors in deposit of, 102
  non-residue for codling moth, 340
  orchard, tar oil, 600
  residual poison, 926
Spreaders, new, for insecticides, 247
Squash bug, control by calcium cyanide, 475
Strawberries, control of white grubs in, 693
Strawberry
  crown borer, poison bait for, 812
  leaf roller, western, control, 121
Sugarcane borer, parasite of, 738
Swallow bug, 949
Sweetpotato
  seed infested by Agromyzid fly, 704
  weevil, control of by methyl bromide, 921

Taeniothrips inconsequens, 950
  simplex, 930
Tar oil
  in delayed dormant sprays, 440
  low temperature, in orchard sprays, 600
Tartar emetic
  against
    onion thrips, 926
    red spider, 579
  toxicity to honeybee, 872
Temperature, effect of on toxicity of stomach poisons, 760
Temperatures
  and insect metabolism, 361
  device for maintaining subzero, 419
Tetranychus telarius, 614
Thermobia domestica, 857
Thrips
  distribution of injurious forms, 133
  on gladiolus, field control, 490
Thrips tabaci, 278, 926
Tick, Gulf Coast, biology of, 179
Tobacco
  flea beetle, 834
    breeding hosts of, 473
    host plants, 467
    insecticides for, 834
    migration of, 470
  moth larvae, control of, 819
  warehouses, fumigation tests in, 812
Tomato fruit worm, control, 129
Tomatoes, control of pentatomids, 275
Toxicity of nicotine and naphthalene, 368
Toxoptera graminum, 628
Traps
  color of, in relation to Japanese beetle, 320
  Japanese beetle, 309
  light
    for leafhoppers, 701
    insect activity at, 654
  value of against Japanese beetle, 317
Treehopper eggs, effect of lime sulfur, and oil, on, 583
Trichopoda pennipes, parasite of *Anasa tristis*, 269
Trout, insect food of, 544
Twig girdlers, 561

Velvetbean caterpillar, control, 635
Veratrum viride extracts, insecticidal action of, 728
Vetch, aphid injury in, 833
Violet, attacked by sawfly, 951
Viruses, plant, and insect vectors, 827
Vitamin content of bee foods, 396

White-fringed beetle, activities of adults, 203
White grubs, control by arsenicals in strawberries, 693
White-marked spider beetle, new common name proposed for, 419
White-wash, to control potato leafhopper, 83
Wireworm, control by
  dichloroethyl ether, 280
  substitute crops, etc., 644

Xylosandrus germanus, 819